

Improving the S/X Celestial Reference Frame in the South

Aletha de Witt¹, Karine Le Bail², Christopher Jacobs³, David Gordon², David Mayer⁴, Matthias Schartner⁴, Sayan Basu¹

Abstract We believe that the S/X celestial reference frame (CRF) can be improved in the far-south by a factor of 2 in density and a factor of 2.5 in precision. We have started a collaboration to meet these goals. We have increased the data rates on existing IVS astrometric sessions in the south from 256 Mbps to 1 Gbps. We will use this sensitivity to detect weaker sources and to improve the precision of sources in the southern S/X CRF, while simultaneously increasing the number of sources, in particular the overlap with other frames such as K- and Ka-band in the radio and the *Gaia* frame in the optical. VLBI observations in the southern celestial hemisphere have always been more difficult both because there are fewer radio telescopes in the south than in the north and because there are fewer known reference sources in the south. There have been many efforts in recent years to increase the number of known reference sources in the south, in particular the LBA calibrator Survey (LCS), which has already produced a significant improvement at X-band. The ICRF-3 is expected to make significant improvements in the south; however, the south has not yet reached parity with the north and much work remains to be done. Therefore, dedicated astrometric and imaging observations have already begun to improve the southern CRF at S/X-bands.

Keywords Astrometry, VLBI, Celestial Reference Frame, Southern Hemisphere, quasars

1. Hartebeesthoek Radio Astronomy Observatory (HartRAO), South Africa
2. NVI, Inc./NASA Goddard Space Flight Center, USA
3. Jet Propulsion Laboratory, California Institute of Technology/NASA, USA
4. Technische Universität Wien, Austria

1 Introduction

Geodetic and astrometric VLBI observations have always been more difficult in the south, with the availability of antennas being the most limiting factor. The second realization of the International Celestial Reference Frame (ICRF-2 [7]) was dominated by data from the north. However, despite many efforts to improve the north/south imbalance of observations (e.g., the AUSTRAL observing program that was started in 2011 [9]), current radio astrometry catalogs are still weak in the south, with a significant hemisphere disparity in source distribution and density.

In recent years there have been many efforts to increase the number of known reference sources in the south, with the most significant contribution coming from the Australian Long Baseline Array (LBA) Calibrator Survey (LCS, [8]), that observed more than 1,500 candidate extragalactic radio sources, (declination below -30°), from 16 VLBI experiments with the LBA at 8.4 GHz.

In 2012, the need for a more uniform spatial coverage of sources and uniform accuracy in source coordinates led to the formation of an International Astronomical Union (IAU) working group, with the goal of the realization of the next generation celestial reference frame (ICRF-3, [6]). Specific emphasis was placed on improving the southern CRF as well extending the frame to higher radio frequencies, chiefly at 24 GHz (K-band [4]) and 32 GHz (Ka-band [5]). Although the ICRF-3 is expected to show significant improvements in the south, the south has not yet reached parity with the north and much work remains to be done.

It is well known that the effect of source structure on astrometric VLBI positions can be significant and that structure and flux density variability are directly

related to the precision of geodetic solutions [3, 10]. It is therefore important to map the structures of these sources on a regular basis. There have, however, only been a few imaging sessions of reference sources in the south and dedicated campaigns to map and monitor source structure have proven difficult to obtain. However, recent investigations to image source structure from existing astrometric and geodetic observations in the south have shown that dedicated imaging campaigns may indeed be possible [1].

In this paper we present the current status of the S/X CRF as well as our proposed plans to improve the S/X CRF in the south. We also present some recent results from these efforts, including a multi-epoch campaign to image source structures in the south.

2 Current Status: North Versus South

At present there are only a few VLBI-capable radio telescopes in the Southern Hemisphere and even fewer that regularly participate in astrometric and geodetic VLBI experiments. The Very Long Baseline Array (VLBA) significantly contributes to CRF work in the north, but unfortunately there is no VLBA to compensate in the south. Currently there are only five radio telescopes in the south that regularly participate in astrometric experiments ($\sim 12\%$ of total); a 15 & 26-m telescope in South Africa, three 12-m telescopes and one 26-m in Australia, and one 12-m telescope in New Zealand. There are two radio telescopes in Antarctica, but these are very small in size and can only detect a few of the brightest sources. There is also a radio telescope in Brazil, but it is close to the equator and does not contribute significantly to southern observations.

In Figure 1 we show the evolution of geodetic and astrometric observations for the period 03 August 1979 to 27 March 2018. The plot shows the distribution of northern-only baselines, southern-only baselines, and north-south or mixed baselines. The distribution has evolved from mainly northern-only baselines to $\sim 10\%$ southern-only and almost 20% mixed baselines.

The growth of astrometric and geodetic observations between the period 03 August 1979 and 27 March 2018 are shown in Figure 2. Southern-only baselines and mixed baselines have increased noticeably in recent years, but still represent only $\sim 15\%$ of the total number of baselines.




Fig. 1 The evolution of the observation distribution from 3 August 1979 to 27 March 2018 between northern-only baselines (blue triangles), southern-only baselines (red inverted triangles), and mixed baselines (green stars).




Fig. 2 Cumulative growth of northern-only observations (blue triangles), southern-only baselines (red inverted triangles), and mixed baseline observations (green stars) for the period 03 August 1979 to 27 March 2018.

The ICRF-2 is based on high precision Very Long Baseline Interferometric (VLBI) measurements of positions of 3,414 extragalactic radio sources. This includes the 295 defining sources which determine the orientation of the frame axes. The ICRF-2 has a noise floor of $40 \mu\text{as}$ in the individual source coordinates, and an axis stability of $10 \mu\text{as}$. The positions were determined from dual-frequency VLBI observations at 2.3 GHz (S-band) and 8.4 GHz (X-band), mostly organized under the auspices of the International VLBI Service for Geodesy and Astrometry (IVS).

The ICRF-2 was generated from 4,726 VLBI sessions and 6.5 million measurements acquired for geodetic and astrometric purposes between 1979 and 2009 and was dominated by data from the north (e.g., Figures 1 & 2). The most recent S/X astrometric so-

lution (sx-gsfc-180521, David Gordon) was generated from 6,206 VLBI sessions and 13.2 million measurements from all available sessions up to 27 March 2018, and includes significantly more southern-only and north-south baseline observations than the ICRF-2.

The sky distribution plot of the formal position uncertainties, from the most recent S/X CRF, is shown in Figure 3. Although this solution shows significant improvement over the ICRF-2, it is clear that we still need more sources in the south and that we also need to improve the spatial coverage, especially for declinations south of -30° . Both the number of sources and the average number of observations per source are a factor of 2 less in the far-south ($\leq -30^\circ$) compared to the far-north. The median formal uncertainties are a factor of 1.5 weaker in $\alpha \cos(\delta)$ in the far-south and a factor of 2.7 weaker in δ . It is evident from these plots that we need more southern baseline observations as well as more north-south baselines.

From the most recent S/X CRF we identified 124 sources (37 ICRF-2 defining sources) in the far-south (below -45° south), with no VLBI images—almost half the total number of sources in the far-south! Multi-epoch maps are essential to assess the astrometric suitability of CRF sources. Extended intrinsic source structures can introduce significant errors in the VLBI measurements, thereby degrading the accuracy of the estimated source positions. The lack of images will severely limit the potential for further improvements in the accuracy of VLBI source positions in the far-south and thus the improved stability of future S/X-band CRFs.

3 Proposed Plans and Progress to Date

3.1 Increase Data Rates

Currently the only dedicated astrometric programs at S/X in the Southern Hemisphere are the IVS Celestial Reference Frame (IVS-CRF) and Celestial Reference Frame Deep South (IVS-CRDS) sessions. Up until 2017, the data rates of these were only 128 Mbps for the IVS-CRF sessions and 256 Mbps for the IVS-CRDS sessions and included only observations of ICRF-2 defining sources. We propose to increase the data rates of these sessions by a factor of 4 or more

by increasing the data rate to 1–2 Gbps. This in turn will allow an increase in the sensitivity by a factor of 2 or more, which will allow the detection of weaker sources down to ~ 350 mJy or less. Scheduling will also become more efficient, since there will be more sources to choose from and scan times will be shorter, which will result in more scans and/or sources per schedule.

We tested and implemented a 1-Gbps observing mode for the IVS-CRDS sessions and a 1-Gbps narrow-band mode was tested for the IVS-CRF sessions. The IVS-CRDS sessions were officially upgraded to 1 Gbps mode starting with crds93 on 24 January 2018. The IVS-CRF sessions switched to 1 Gbps on 4 April 2018 with crf106.

3.2 Scheduling Optimized for Astrometry

We propose to change the scheduling of the IVS-CRF and IVS-CRDS sessions to be optimized for astrometry and imaging instead of geodesy. This would imply using the full network of stations when possible for every scan and with no sub-netting as is used routinely for geodesy schedules. There should also be at least 3–8 scans per source spread evenly over hour angle range, to allow for optimal u - v coverage for imaging without compromising the astrometric goals of the experiment. The schedule should also include blocks with tropospheric calibrators, that will also be used as astrometric ties and for amplitude calibration for imaging. In addition, we propose astrometric sessions be scheduled as part of a campaign rather than individual sessions. This will ensure that each source will receive the required amount of observing time and that the ultimate astrometric goals of the project be reached.

We have optimized the scheduling of all of the IVS-CRDS sessions from crds93 onwards. Figure 4 compares the u - v coverage for a source observed in both crds68 (27 November 2013) and crds94 (21 March 2018). The improvement in u - v coverage going from two scans (crds68) to seven scans (crds94) is clearly evident from these two plots. The overall number of sources also increased from 38 to 51 and the overall number of scans from 144 to 304, from crds68 to crds94.




Fig. 3 The distribution of sources from the most recent S/X astrometric solution (sx-gsfc-180521) showing the formal uncertainties in $\alpha \cos(\delta)$ on the left and δ on the right.




Fig. 4 The u - v plane coverage for the source 1420-679, observed in different IVS-CRDS sessions. The u - v coverage plot on the left is from the crds68 session (27 November 2013) and the plot on the right, showing a much improved sampling of the u - v plane, is from the crds94 session (21 March 2018). In both sessions five antennas participated. In crds68 the source was only observed in two scans and in the crds94 it was observed in seven scans.

3.3 Improve Precision

Up to 2017, only ICRF-2 defining sources were observed in IVS-CRF and IVS-CRDS sessions. We propose to re-observe all southern sources in the current S/X CRF to improve the source position accuracy in both coordinates. We propose to improve the overall precision by a factor of 2.5 in the south. From the 1,344 sources south of -15° , we have 1,091 sources with ≤ 10 observing sessions. We will prioritize the 216 of these sources with flux density > 350 mJy that will be easily detectable with current instruments and data rates. Since December 2017, we started to include some of these 216 sources as part of the IVS-CRDS and IVS-CRF sessions.

3.4 Improve Density and Spatial Coverage

We propose to improve the far-south by a factor of 2 in density by expanding the source list in the south, specifically in the far-south (below -30° south). In addition, we also propose to improve the overlap with K- and Ka-band frames and the *Gaia* optical frame. We identified ~ 80 K- and Ka-band sources that are not in the current S/X frame at declinations south of -15° . From these we have ~ 20 sources with flux density > 350 mJy at S/X-band. In addition, we also propose follow-up observations of candidate CRF sources brighter than 350 mJy from the pool of LCS sources. Priority will be given to $\sim 1/2$ of the target sources that have a counterpart with *Gaia*. Since December 2017, we started to include those K- and Ka-band sources that are not currently in the S/X CRF into the IVS-CRDS sessions.

3.5 Imaging

We propose to produce multi-epoch maps at both 2.3 and 8.4 GHz for all sources observed in the IVS-CRDS and IVS-CRF sessions. These maps will be used to quantify the non-pointlike structure and jet directions in these CRF sources. First priority will be given to those 124 sources in the far-south with no VLBI images. In Figure 5 we show representative contour plots from recent imaging results obtained from the crds94 session from 21 March 2018 [2].




Fig. 5 From left to right, contour plots for sources 1045-620, 1646-506, 1657-582, and 1831-711 at 8.6 GHz from 21 March 2018 (crds94). North is Up and East is to the Left. The FWHM beamsize is graphically indicated in the bottom left corner [2].

4 Conclusions

Our goal is to improve the S/X-band frame in the south by at least a factor of 2 in density and 2.5 in precision, to be about as good as the north.

In order to achieve this, we propose to increase the data rate of southern IVS sessions to at least 1 Gbps and to optimize the scheduling of these sessions for astrometry and imaging versus geodesy. We further propose to increase the number of well observed sources ($N_{\text{sess}} > 10$) in the south and to increase both the number of southern-only and north-south baseline observations. We also propose to expand the southern source list and improve spatial coverage. In addition, we propose multi-epoch imaging of southern CRF sources to quantify non-pointlike structure and measure jet directions.

Our initial steps are succeeding: all IVS southern astrometric sessions are now at 1 Gbps, non-defining sources were added to the IVS-CRDS and IVS-CRF source list, and we produced first imaging results from and IVS-CRDS sessions at 1 Gbps.

Acknowledgements

Copyright 2018. All rights reserved. U.S. Government sponsorship acknowledged for part of this research. HartRAO is a facility of the National Research Foundation (NRF) of South Africa. The Hobart telescope is operated by the University of Tasmania and this research has been supported by AuScope Ltd., funded under the National Collaborative Research Infrastructure Strategy (NCRIS).

References

1. S. Basu et al., How Good is the Deep Southern Sky, in D. Behrend, K. D. Baver, K. L. Armstrong (eds.), *IVS 2016 General Meeting Proceedings: "New Horizons with VGOS"*, p. 312–316, 2016.
2. S. Basu et al., VLBI Imaging Observations of Potential ICRF-3 Defining Sources in the South. *MNRAS*, in preparation, 2018.
3. P. Charlot., Radio-source Structure in Astrometric and Geodetic Very Long Baseline Interferometry, *AJ*, **99**, 1309, 1990.
4. A. de Witt et al., K-band Celestial Reference Frame: Can it be Better than S/X?, in R. Haas & G. Elgerd (eds.), *Proceedings of the 23rd EVGA Working Meeting, May 2017, Gothenburg, Sweden*, pp. 181–185, 2017.
5. C. Jacobs et al., Celestial Reference Frame at X/KA-Band (8.4/32 GHz) for Deep Space Navigation, *23rd International Symposium on Space Flight Dynamics, Pasadena CA, 30 Oct 2012*, 14 pages, id.1, 2012
6. C. Jacobs et al., ICRF-3: Roadmap to the Next Generation ICRF, in N. Capitaine (eds.), *Proceedings of the Journées 2013, 16-18 September 2013*, p. 51–56, 2014.
7. C. Ma et al., The Second Realization of the International Celestial Reference Frame by Very Long Baseline Interferometry, *IERS Technical Note 35*, 29, 2009.
8. L. Petrov et al., The LBA Calibrator Survey of Southern Compact Extragalactic Radio Sources - LCS1, *MNRAS*, **414**, 2528, 2011
9. L. Plank et al., The AUSTRAL VLBI Observing Program, *Journal of Geodesy*, **91**, 7, p. 803–817, 2017.
10. S. Shabala et al., Quasar Structure Effects on the VLBI Reference Frame: The Case of 1144-379, In D. Behrend and K.D. Baver (eds.), *Seventh IVS General Meeting*, 329, 2012.